



Science Mission Directorate
National Aeronautics and Space Administration

UAH

AIRS DATA ASSIMILATION AT THE SPoRT CENTER

Brad Zavodsky

University of Alabama – Huntsville
Huntsville, Alabama

and


Gary Jedlovec & Shih-hung Chou

NASA / Marshall Space Flight Center
Huntsville, Alabama

AIRS Science Team Meeting – March 2006



transitioning unique NASA data and research technologies to the NWS



Presentation Outline

- SPoRT overview
- Recap of SPoRT AIRS profile assimilation results presented at September 2005 AIRS Science Team Meeting
- New case study (w/ v4.7 profile assimilation)—impact on analysis and forecast and comparison with v4.0
- Future plans with AIRS profiles and radiances





NASA's Short-term Prediction and Research Transition (SPoRT) Center

Mission: Apply NASA measurement systems and unique Earth science research to improve the accuracy of short-term (0-24 hr) weather prediction at the regional and local scale (<http://weather.msfc.nasa.gov/sport/>)

Test-bed for rapid prototyping of new products

Transition research capabilities / products to operations

- real-time MODIS data and products to 6 NWS forecast offices
- twice daily WRF model output (initialized with MODIS SSTs)- operational
- convective initiation / lightning products for nowcasting severe weather

Development of new products and capabilities for transition

- MODIS SST composites, AMSR-E rain rates, and ocean color products
- assimilation of AIRS radiances and thermodynamic profiles into regional
- forecast models



transitioning unique NASA data and research technologies to the NWS




Motivation / Relevance of SPoRT

SPoRT activities are problem driven

- apply Earth science technologies to regional weather studies

Focus on:

- nowcasting – lightning, CI, diagnose visibility and ceiling for aviation
- advanced modeling and data assimilation – MODIS and AIRS in WRF – regional as opposed to global applications addressed by JSCDA / GMAO
- MODIS and AMSR-E data and products



Relevant to several key NASA Earth science questions

- improvements in weather forecasting
- regional air quality impacts
- relationship between weather and climate variability

Direct connections to areas on National Applications

- water and air quality
- disaster coastal management
- aviation



transitioning unique NASA data and research technologies to the NWS




AIRS Profile Assimilation in Regional Models

*Straight-forward way to have
AIRS affect forecasts*

AIRS Science Team Meeting (Sept. 2005)



- described use of QIs in v4.0 in the ADAS/WRF system
- AIRS v4.0 retrievals had a positive impact on 0-48h regional forecast for west coast U.S. case study
 - treat AIRS profiles similar to RAOB
 - tune ADAS 3D influence parameters for AIRS
 - use QIs to eliminate some profiles and to assign weight of data influence
- need for more (refined) quality indicators
 - moisture QIs separate from temperature
 - better vertical delineation in quality

AIRS assimilated at 2200UTC on 14 January 2004




ADAS Background

BKGD+AIRS+MADIS



AIRS analysis

Impact of DA



Example error profile for ADAS for AIRS data failing low-level temperature check






New Case Study: November 20-22, 2005


Rapidly intensifying storm off the eastern seaboard poorly forecasted by GFS, NAM, and SPO_{RT} operational WRF

Case Selection

- relevant to SPO_{RT} interests in SEUS region
- more verification data available over the Eastern US (minimal terrain impact)
- synoptic setting
- opportunity to eventually test both over-ocean and over-land AIRS profiles (current case study focuses solely on over-ocean retrievals)
- comparable CONUS domain to operational SPO_{RT} WRF for easy transfer to operations




Surface analysis 11/20/05 12Z



Surface analysis 11/22/05 12Z



transitioning unique NASA data and research technologies to the NWS



AIRS Data Overview


QI improvements in v4.7:

- pressure for each sounding indicating level of valid data
- level-by-level error estimates for each T and q profile


More data are assimilated

- number of assimilated profiles reduced in v4.7
- however, higher data volume as more data are used in the mid-troposphere (previously ambiguous QI here)
- assimilating a larger volume of higher quality data should produce an analysis that provides better initial conditions for the WRF

Version 4.0



Version 4.7



Let's take a closer look at the differences between the v4.0 and v4.7 temperature and moisture profiles






963 AIRS profiles assimilated 797 AIRS profiles assimilated



transitioning unique NASA data and research technologies to the NWS



v4.0/v4.7 AIRS Profile Comparison



30.512°N, 72.342°W (RED circle)

38.961°N, 72.389°W (BLUE circle)

24.364°N, 69.991°W (GREEN circle)

Black: v4.0 (standard; 12 levels) Red: v4.7 (supplemental; ≈ 53 levels)

- Small change in T but significant q reduction (drying) in the mid-troposphere (evident even with different vertical resolution)
- Changes made to the water vapor regression step



transitioning unique NASA data and research technologies to the NWS




SPoRT Research WRF and Analysis

WRF Configuration

- 36km domain with 37 vertical levels
- state-of-the-art dynamics and physics
- initialized with NCEP 1° GFS grids, with 6-h forecasts used as LBC


ADAS Configuration

- same resolution as WRF (43 vertical levels)
- Bratseth SCM weights obs using horizontal and vertical ROIs and error characteristics
- error table for each data: AIRS errors are from v4.0 documentation ($\approx 1/3$ of BKGD)
- sufficient vertical scaling to account for the difference in vertical resolution of AIRS soundings




Assimilation / Forecast


- GFS interpolated to WRF domain
- WRF analysis at 18 UTC used as background for ADAS



transitioning unique NASA data and research technologies to the NWS









v4.0



WRF Background
at 18 UTC

v4.7




- significant analysis/background differences ($\approx 20\%$ of total q at 700 mb)
- v4.7 analysis is ≈ 0.5 g/kg drier than the v4.0 throughout most of the AIRS swath
- difference between v4.7 and v4.0 impact on temperature analysis field generally smaller






30h Forecast Validation Against RAOBs



Validated against
17 east coast
RAOBs




- inclusion of v4.7 AIRS data reduces bias in T and q at most levels
- RMS error is smaller for T below 600 mb with v4.7 data; control performs better above 600 mb
- RMS error improvements for q are mixed with AIRS improving forecast in mid-troposphere





30h Forecast Intercomparison

700 mb T and q



v4.7 vs. v4.0 forecast verification impact on RMSE greater for T but q significantly impacts forecast outside validation region

- $\pm 1^\circ$ temperature differences over many parts of coastline where verification occurs
- largest differences (> 1 g/kg) in moisture field occur outside verification region



transitioning unique NASA data and research technologies to the NWS



November 2005 Case Study Summary

New level-specific QIs for v4.7 provide a larger volume of higher quality data for assimilation


v4.7 data improve the 30h forecast of T and q compared to RAOBs

- forecast bias improves over control and v4.0 for both T and q at most levels
- forecast RMS significantly improves over the control and v4.0 for T below 600 mb; RMS results are mixed for q in validation region
- improvements are likely due to improved moisture regression and more specific QIs in the v4.7 data

Future plans involving AIRS

- developing case-specific error tables for ADAS using a statistical representation of the individual profile error estimates in the new v4.7 data
- assimilation of AIRS radiance data on regional scale





Radiance Data Assimilation into Regional Models

Direct way to have AIRS data affect short term weather forecasts

Use state-of-the-art WRF modeling system


- future NCEP operational system
- data assimilation using Gridpoint Statistical Interpolation (GSI) – 3DVAR

GSI supports cloud-free radiance data assimilation

- more direct method than profiles – avoids retrieval problems
- single field of view data
- existing approach with GOES / TOVS
- consider cloud-cleared radiances – quality? - data volume?

Problems with this approach:

- data volume – large even on a regional scale
 - “super obs” - brute force, not necessarily the best
 - need intelligent data thinning – based on weather of the day
- need to identify cloud-free radiances
 - based on channel not cloud mask







Detection of Clear / Cloudy Radiances

Cloud mask used to identify clear-cloudy AIRS ifovs

- if cloudy throw away all radiance data
- 5-10% clear AIRS ifovs – usually not meteorologically significant regions
- need to identify cloud contamination by channel



Use CO2 sorting approach to explicitly identify channels contaminated by clouds

- sort data by brightness temperature
- determination of the separation point between contaminated and uncontaminated channels
 - position of the separation point is a function of CTP
 - magnitude of the separation is a function of ECF
- clear scene is dependant on a background field

Impact:

- 2-3 factor increase in radiances (over masking approach)
- Data added in meteorologically significant regions (above clouds)





Towards Operational Assimilation of AIRS Data on a Regional Scale

Develop operational capability for real-time AIRS data assimilation as supplemental forecasts for NWS WFOs in Southeast U.S.

- ADAS/WRF with AIRS profiles in FY06-FY07
- GSI / WRF – cloud-free radiances in FY07-FY08

Considerations:

- real-time AIRS availability
- sensor web capabilities for autonomous operations—select forecast/assimilation cycle based on:
 - anticipated weather (high impact events)
 - juxtaposition of AIRS pass w.r.t. the storm feature of interest
 - other ancillary data
- intelligent data thinning

